

<https://helda.helsinki.fi>

Using the first European Breeding Bird Atlas for science and perspectives for the new Atlas

Herrando, Sergi

2019

Herrando , S , Keller , V , Bauer , H-G , Brotons , L , Eaton , M , Kalyakin , M , Voltzit , O ,
Lehikoinen , A , Milanesi , P , Noble , D , Ramirez , I , Vorisek , P & Foppen , R 2019 , '
Using the first European Breeding Bird Atlas for science and perspectives for the new Atlas '
, Bird Study , vol. 66 , no. 2 , pp. 149-158 . <https://doi.org/10.1080/00063657.2019.1618242>

<http://hdl.handle.net/10138/312708>

<https://doi.org/10.1080/00063657.2019.1618242>

unspecified

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Using the first European Breeding Bird Atlas for science and perspectives for the new Atlas

SERGI HERRANDO^{1,2}, VERENA KELLER^{1,3}, HANS-GÜNTHER BAUER^{1,4}, LLUÍS BROTONS^{1,2,5,6}, MARK EATON^{1,7}, MIKHAIL KALYAKIN^{1,8}, OLGA VOLTZIT⁸, ALEKSI LEHIKONEN^{1,9}, PIETRO MILANESI^{1,3}, DAVID NOBLE^{1,10}, IVÁN RAMÍREZ^{1,11} PETR VOŘÍŠEK^{1,12} & RUUD FOPPEN^{1,13,14}

1 European Bird Census Council, PO Box 6521, 6503 GA Nijmegen, The Netherlands.

2 Catalan Ornithological Institute. Natural History Museum of Barcelona. Pl. Leonardo da Vinci 4-5. 08019 Barcelona, Spain.

3 Swiss Ornithological Institute. Seerose 1. CH-6204 Sempach, Switzerland.

4 Max-Planck-Institut für Ornithologie, Vogelwarte Radolfzell, Am Obstberg 1, 78315 Radolfzell Germany.

5 InForest Jru (CTFC-CREAF), Crta. Antiga St Llorenç de Morunys km 2, 25280, Solsona, Spain.

6 CSIC, Cerdanyola del Vallès, Spain.

7 The RSPB Centre for Conservation Science, RSPB, Sirius House, 1 Amethyst Road, Newcastle Business Park, Newcastle-upon-Tyne, United Kingdom.

8 Zoological Museum of Lomonosov Moscow State University, Bolshaya Nikitskaya Str., 2, Moscow, 125009, Russia.

9 The Helsinki Lab of Ornithology, Finnish Museum of Natural History, University of Helsinki, Finland.

10 British Trust for Ornithology, The Nunnery, Thetford, Norfolk IP24 2PU, UK.

11 BirdLife International, David Attenborough Building, Pembroke Street, Cambridge, CB2 3QZ, UK.

25 12 Czech Society for Ornithology. Na Belidle 34 CZ-150 00 Prague 5, Czech Republic.

26 13 Sovon Dutch Centre for Field Ornithology, PO Box 6521, 6503 GA Nijmegen, The

27 Netherlands.

28 14 Department of Animal Ecology and Ecophysiology, Institute for Water and Wetland Research,

29 Radboud University, 6500, Nijmegen, The Netherlands.

30

31 **SHORT TITLE:** The European Bird Atlas and science

32

33 **KEYWORDS:** bird species ranges, European Bird Census Council, Western Palearctic,

34 bibliographic research.

35

36 **EMAIL ADDRESS OF CORRESPONDENCE AUTHOR:** ornitologia@ornitologia.org

37

38

38

39 **SUMMARY**

40 **Capsule** The first EBCC Atlas of European Breeding Birds (EBBA1) has been widely
41 used in scientific publications.

42 **Aims** To quantify how scientific publications have used data from EBBA1, what the
43 topics of these studies have been, and to identify key aspects in which the Second
44 European Breeding Bird Atlas (EBBA2) will provide new opportunities for basic and
45 applied science.

46 **Methods** We searched for “EBCC Atlas of European Breeding Birds” in Google
47 Scholar to find papers published in scientific journals that cited EBBA1. We analysed
48 the contents of a random selection of 100 papers citing this atlas. We described the
49 way these papers used EBBA1.

50 **Results** EBBA1 has been cited in 3 150 scientific publications, and can be regarded as
51 a fundamental reference for studies about birds in Europe. EBBA1 was extensively
52 used as a key reference for the studied bird species. A substantial number of papers
53 re-analysed EBBA1 data to derive new information on species distribution, ecological
54 traits and population sizes. Distribution and ecology were the most frequent topics of
55 studies referring to EBBA1, but this source of information was employed in a very
56 diverse range of studies. In this context, climate change, impact of agriculture and
57 habitat loss were, by order, the most frequently studied environmental pressures.
58 Constraints in EBBA1 such as the poor coverage in the East of Europe, the lack of
59 information on distribution change and the coarse resolution were identified as issues
60 limiting the use of EBBA1 for some purposes.

61 **Conclusions** This study demonstrates the scientific value of European-wide breeding
62 bird atlases. EBBA2, with its almost complete coverage across Europe, the
63 incorporation of changes in distribution between the two atlases and the inclusion of

modelled maps at a resolution of 10x10 km will certainly become a key data source and reference for researchers in the near future.

INTRODUCTION

The collection of biodiversity data over large geographical areas is essential in order to adequately inform conservation policy (Chiarucci *et al.* 2011, Schmeller *et al.* 2015). Great efforts have been made in recent years to integrate biodiversity information from thousands of sources into global data depositories, such as the Global Biodiversity Information Facility (GBIF 2019) and the Map of Life (MOL 2019), among others. These depositories gather a huge amount of open access data, and thus allow scientists to address relevant issues on biodiversity and conservation. However, an aspect that is sometimes neglected is that differences with respect to data sampling, storage and mobilization often lead to bias in large-scale biodiversity patterns (Beck *et al.* 2014).

Biodiversity data are compiled in very different ways across Europe (Voříšek & Marchant 2003, EuMon 2019). In Western and Central European countries, citizen science - understood as a strong partnership between amateur and professional scientists - has constituted a fundamental pillar for the compilation of biodiversity data (Bell *et al.* 2008, Dickinson *et al.* 2010). In contrast, the tradition of data collection in the eastern half of Europe and the Balkans has been often based on professional work directed by academies of science or similar governmental institutions (e.g. Budinski *et al.* 2010). In addition, methodological differences among countries are influenced by socio-political contexts, specific national targets, fieldwork and data mobilization approaches, and geographical particularities.

Conscious of the value of integrating national data to aid the understanding of large-scale patterns of distribution and abundance in birds, European ornithologists have a long tradition of cooperation among national organizations. Developing from a series of meetings aimed at greater international collaboration in ornithology, the European

Ornithological Atlas Committee (EOAC) was set up in 1972. Its representatives soon realized that a Europe-wide atlas could be produced from national surveys if common standards were adopted (see Greenwood 2017). The EOAC later changed its name and formalized its organizational structure to become the European Bird Census Council (EBCC), the organization that produced the EBCC Atlas of European Breeding Birds (Hagemeijer & Blair 1997), hereafter named EBBA1.

EBBA1 integrated data on the distribution and abundance of breeding birds, compiled at national level, into a single pan-European collaborative work. Thanks to the coordinated efforts of thousands of ornithologists and birdwatchers, accurate information on distribution and abundance was gathered for 495 species in 3 950 50x50 km squares (Figure 1). After publication EBBA1 data were made accessible to researchers, first under direct request to the EBCC Executive Committee and, since 2015, as open access via GBIF. EBBA1 has represented one of the few sources of information concerning distribution of bird populations at a continental scale and consequently has been widely employed in fundamental and applied research and for conservation-oriented publications (e.g. BirdLife International 2004). However, an analysis of the use of EBBA1 in research to assess its impact in the scientific community has not been done.

In this paper we show the potential of using such pan-European distribution data in research. More specifically, we review how EBBA1 was used in scientific literature and what the aims of these studies were. At the moment of writing this article, the Second European Breeding Bird Atlas (EBBA2) (Keller 2014, EBBA2 2019) is finalizing the compilation of updated data on the distribution of all breeding bird species (Figure 1). Following this review of literature we discuss how this knowledge has been incorporated in the aims, organization and methodological choices for the EBBA2 project. Finally, we show our vision of current efforts in the integration of continental bird data and anticipate how EBBA2 will contribute to basic and applied research.

118

119 MATERIALS AND METHODS

120 We considered Google Scholar as our source of information to analyse the use of
121 EBBA1 in scientific publications. This search facility is freely and easily accessible,
122 provides similar results to Thompson's ISI Citation Index (Pauly & Stergiou 2005), and
123 it has become increasingly popular in recent years (de Winter *et al.* 2014). We
124 conducted the literature search in December 2016. We queried the sequence of words
125 'EBCC Atlas of European Breeding Birds' and downloaded results in a database.
126 Despite the potential scientific interest of information collected in books, theses,
127 technical reviews and 'grey' literature, we removed these types of works from the
128 original database and kept only papers published in scientific journals. Then we
129 performed a random selection of 100 papers from the total available using the
130 Wichmann–Hill number generator (Wichmann & Hill 1982). Selected papers (shown
131 with a symbol * in this article) were read by SH to analyse the general use of EBBA1 in
132 science.

133 We addressed three specific questions related to how EBBA1 was used in scientific
134 publications:

- 135 • Was EBBA1 used only for its information about birds? Or did it also attract
136 researchers' attention regarding the way the project was organised?
- 137 • Were EBBA1 data re-analysed in these scientific publications?
- 138 • When new analyses of EBBA1 were performed, what were the purposes?

139 In addition, we addressed five questions related to the subject matter of the papers that
140 cited EBBA1:

- 141 • What were the major research topics of papers that cited EBBA1?

- What were the pressures on biodiversity addressed in the papers that cited EBBA1?
- Was EBBA1 used more often in papers that studied observed change in the past or predicted future change?
- Was EBBA1 more often used in international, national or subnational studies?
- Was EBBA1 more often used in studies about a single bird species, multiple bird species or studies that included several groups (birds plus others)?

RESULTS

How EBBA1 was used in scientific publications

Our search in Google Scholar showed that 3 150 scientific papers published in journals referred to 'EBCC Atlas of European Breeding Birds'. From the 100 papers randomly selected for this study 87 papers were written in English and 13 in nine other languages: German (5), Russian (1), Spanish (1), French (1), Italian (1), Polish (1), Dutch (1), Catalan (1) and Norwegian (1). All these papers are listed in Appendix I.

A total of 96 papers analysed in this study were interested in the EBBA1 data on breeding bird species and four papers focused on the methodology and on how the project was organized. A total of 76 papers referred to the general patterns of species distribution, abundance or ecology shown in EBBA1, but did not use atlas data to conduct new analyses. Nevertheless, 19 papers re-analysed EBBA1 data for three main purposes: analyses of species distribution (8 papers), determining species ecological traits (7) and analyses focusing on population sizes (4).

Subject matter of the papers that cited EBBA1

Main topics

EBBA1 was cited for many different scientific purposes within the field of biodiversity research. Nevertheless, two topics stood out: species distribution (50 papers) and

ecology (44 papers). Other topics such as climate, conservation, trends or abundance were addressed by a smaller number of papers (Figure 2).

Pressures on biodiversity

Climate change was the environmental pressure more frequently reported in papers referencing EBBA1, accounting for a total of 12 papers. Concern on how changes in agricultural practices are affecting farmland birds was the topic of five papers, habitat loss and environmental impact associated with invasive alien species were studied in four papers each and pollution was addressed in three papers.

Temporal focus

EBBA1 was a reference for 22 papers that incorporated an assessment of temporal change in past-to-present approaches. Future projections were at least partly addressed in five papers.

Scale

Our survey showed that 54 of the analysed papers studied patterns at subnational level, 28 papers at national level; 18 papers worked at the European scale (or substantial part of the continent) and five addressed research at the global scale.

Spatial resolution

EBBA1 presented the data in a relatively coarse 50x50 km grid. Among the studies citing EBBA1, 16 studied biogeographical or macroecological patterns at the same spatial resolution used in this atlas. In 25 papers analyses were conducted at finer resolutions, usually using EBBA1 as background information but in few cases re-analysing its data and then applying the results at finer resolution.

Biodiversity focus

EBBA1 was cited predominantly in papers on birds, namely in 48 papers on single species and 46 treating several species. Five papers focused on birds and other biological groups. Only one paper was not interested in birds but exclusively in plants.

DISCUSSION

How EBBA1 was used in scientific publications

Our analyses show that EBBA1 was used in very different ways. Very often EBBA1 was employed as background information for studies about a particular species, such as that of the status of the Bearded Parrotbill *Panurus biarmicus* in Italy (Brichetti & Grattini 2008)* or the decline of the Red-backed Shrike *Lanius collurio* in Zealand, Denmark (Pedersen *et al.* 2011)*. In other cases, EBBA1 information was viewed in a wider context and represented a general reference for the distribution of groups of bird species, such as for the ranges of boreal birds (Virkkala & Rajasärkkä 2007)*.

A substantial proportion of the sampled papers not only cited EBBA1 data but presented novel analyses using it. Some of the papers that re-analysed the data were interested in deriving information on species distribution. For example, EBBA1 data were used (together with data from the Atlas of Southern African Birds, Harrison *et al.* 1997), to model species distributions at the Last Glacial Maximum and then analyse the contrasting late-Quaternary histories for Southern and Northern Hemisphere bird species (Huntley *et al.* 2013)*. EBBA1 was also used to derive species ecological traits based on the species' respective distributions. For example Gaüzère *et al.* (2015)* used EBBA1 data to calculate species-specific temperature indexes (STI), representing the preferred climate conditions of species based on temperatures within their area of distribution and then showed a fine-scale and short-term adjustment of community composition to temperature changes. An example of EBBA1 data use to derive information on species population size is the study of Møller *et al.* (2011)* who

215 obtained population sizes from this atlas to determine that migratory divides constitute
216 barriers to dispersal with consequences for ecology and evolution of populations.

217 **Topics in papers referring to EBBA1**

218 *Main topics*

219 EBBA1 was widely used as a reference for bird species distribution, population size
220 and ecology in many scientific publications interested in a variety of scientific topics.
221 Many distributional studies conducted at national or regional scales referred to EBBA1
222 to provide a wide geographical context for the species of interest. This background
223 information was applied in a variety of circumstances, such as the delimitation of
224 Important Bird Areas (IBAs) in Macedonia (Veleviski *et al.* 2010)* and the distribution of
225 birds of prey in Belarus (Dombrovski & Ivanovski 2005)*. Information from EBBA1 was
226 widely used as a suitable reference in papers on species ecology. A first example is a
227 study that used bird-land cover associations based on EBBA1 data in order to
228 determine the influence of this predictor (and others) on beta diversity for European
229 birds (Keil *et al.* 2012)*. A second example is the use of EBBA1 as a reference for the
230 habitat preferences of Dartford Warbler *Sylvia undata* in the context of assessing its
231 status in the UK (Wotton *et al.* 2009)*.

232 Conservation also emerged as a very relevant topic and information from EBBA1 was
233 used for a wide variety of purposes. For instance, the evaluation of the network of
234 protected areas by Sándor & Domşa (2012)* who determined additional areas to better
235 preserve forest specialist birds on the basis of a thorough analysis of habitat
236 preferences, the assessment of species of conservation concern as described by Keller
237 *et al.* (2010)* who assessed the international importance of the breeding populations of
238 bird species in Switzerland by estimating the percentage of the national population
239 using EBBA1 as reference, and the evaluation of land use policies as described by

Eggers *et al.* (2009)*, who used the EBBA1 dataset to evaluate whether biofuel policies were detrimental to biodiversity in agricultural landscapes in Europe.

Pressures on biodiversity

Five out of the six main categories of environmental pressure according to Aichi Target B (Secretariat of the Convention on Biological Diversity 2014) were found as topics in the analysed papers, namely climate change, agricultural practices, habitat loss, invasive alien species and pollution. EBBA1 data had a prominent role in modelling climatic ranges of bird species in the context of predictions of impact of climate change on their populations either at continental (e.g. Huntley *et al.* 2004)*, national (e.g. Reif *et al.* 2013, Koleček *et al.* 2014)* and local level (e.g. Flousek *et al.* 2015)*. Beyond our random selection of papers in journals, it is noticeable to mention that EBBA1 was essential to model European species distributions under future climatic scenarios (Huntley *et al.* 2008). Effects of agricultural practices on birds was also addressed by some of the analysed papers. For instance, Meller *et al.* (2015)* used EBBA1 data and predicted that the magnitude of range shifts due to climate change was far greater than the impact of land conversion to bioenergy plantations within the European Union.

Temporal focus

Since EBBA1 maps provided just a single time reference, and there is no comparable data source for another period yet, no study used the data from this atlas to explicitly analyse temporal changes in bird distribution. However, some authors used the information written in the species accounts in EBBA1 to describe general patterns of distribution or population change, such as in the study of the decline in Tree Sparrow *Passer montanus* populations in Poland (Tomiałoć 2012)*. EBBA1 data were cited in several publications to project species distributions into the future, such as the prediction of a considerable risk of extinction within the Bavarian Forest National Park as a result of global warming (Bässler *et al.* 2010).

266 *Scale*

267 To date EBBA1 is certainly among the most relevant original works that map the
268 occurrence of breeding birds in the whole of Europe. Therefore, it has been widely
269 employed as a key reference for the European distribution of birds in studies carried
270 out at subnational, national, continental and global scales. For example, Chiron *et al.*
271 (2009)* used EBBA1 data to explore the relationship between the geographical pattern
272 of richness of native bird species and that of exotic bird species in Europe. However,
273 EBBA1 was mostly used as a reference concerning species distribution at subnational
274 level, such as in the study of the impact of radiation from the Chernobyl Nuclear Power
275 Plant on birds by Møller *et al.* (2010)*.

276 *Spatial resolution*

277 The detail at which a geographical dataset defines the location of its features is a key
278 element for studies on species distribution. EBBA1 provided data in a 50x50 km grid
279 and, as shown in many of the examples mentioned so far, this cell size met the needs
280 of many of the analyses required to make progress in diverse fields of the ornithology.
281 For example, Carrascal & Seoane (2009)* aggregated original 10x10 km data for
282 Bonelli's Eagle *Aquila fasciata* from a Spanish atlas to a coarser 50x50 km resolution
283 because they considered that in the context of analysing of factors affecting its
284 distribution in Spain, the latter provided a higher certainty about the species occurrence
285 in each cell (absences more likely to be real).

286 *Biodiversity focus*

287 EBBA1 has undoubtedly created one of the most complete databases on the
288 distribution of birds in Europe. Other large-scale studies have reached a similar status
289 for other taxonomic groups. Together they have conformed an excellent dataset for
290 many studies interested in biodiversity in a broader sense. For example, Huntley *et al.*

291 (2004)* used distribution data from plants, birds and insects to predict that climatic
292 model performance was related neither to major taxonomic group nor to trophic level.
293

294 **PERSPECTIVES FOR THE NEW EUROPEAN BREEDING BIRD ATLAS**

295 At the moment of compiling information for the analyses shown in this article, a total of
296 3 150 scientific journal papers referred to the 'EBCC Atlas of European Breeding Birds'.
297 A new Google Scholar search would find that this is clearly an outdated number. All
298 these values illustrate the recognition EBBA1 has received in the scientific community,
299 but also the very high research potential of an atlas update.

300 The analysis of publications citing EBBA1 revealed issues that are important to be
301 taken into account for other projects and in particular for the Second European
302 Breeding Bird Atlas, hereafter called EBBA2. Some of the papers analysed in this
303 study made specific comments on the limitations of EBBA1 data, referring especially to
304 deficiencies in the coverage, such as in the eastern part of the continent and the
305 Balkans (e.g. Keil *et al.* 2012)*. EBBA1 gaps of information resulted mainly from
306 limitations in coordination, fieldwork capacity and data flow at that time. To launch the
307 new project and update in the best possible manner the EBBA1 information, EBCC
308 constituted an EBBA2 steering committee whose members had experience in
309 organising surveys, analysing the data and interpreting the results. These members
310 came from different parts of Europe because bird atlas approaches differ depending on
311 the biogeographical, cultural, traditional and economic contexts (Greenwood 2007).
312 Then, an EBBA2 coordination team was established to deliver the central functions of
313 running the atlas project and of keeping a fluent communication within the network of
314 scientists developing methodologies at the European level, coordinators organising
315 fieldwork at national level and skilled ornithologists doing the fieldwork. This close

collaboration guaranteed the top-down integration from a bottom-up approach taking the very diverse situation in Europe into account (Herrando *et al.* 2014).

The analysis of the papers that cited EBBA1 showed that many studies focused on changes in occurrence. However, none could use that dataset to study changes in distribution (see *Temporal focus*). Updating EBBA1 information will provide many opportunities to analyse changes in distribution and, thus to investigate the environmental pressures driving these changes. Therefore, one of the most important methodological aspects of EBBA2 was to maximize comparability between the two atlases. Consequently, the same UTM 50x50 km grid, breeding codes and abundance codes as used in EBBA1 were adopted for EBBA2. This grid was also selected in compliance with projects working on other taxonomic groups, e.g. see Sillero *et al.* (2014) for the new atlas of amphibians and reptiles of Europe, which will provide a good basis for cross-taxa analyses in the future (see *Biodiversity focus*). However, robust analyses of change do not only depend on the use of the same geographical units but also on the particularities of the fieldwork done in each square in each atlas. To overcome this difficulty, EBBA1-EBBA2 species distribution change was incorporated in the EBBA2 project by means of taking into account information on the fieldwork process for the two atlases. In fact, change was not calculated for all squares but for a subsample that were sufficiently well covered in the two atlases. These squares were determined after a series of analyses that incorporated information on the reported completeness of coverage and the total number of species detected as well as their deviance from modelled expected richness, among others. Finally, in the overall framework of improving the quality of the data, EBBA2 implemented a novel approach by incorporating highly standardised surveys within a sample of 10x10 km squares. The more immediate aim of this survey was to generate detailed maps of species breeding occurrence (see next paragraph), but these standardised surveys

could provide potential added value for robust comparisons in the future (see approach implemented for this purpose in Balmer *et al.* 2013).

Gaston *et al.* (2008)* pointed out that, despite the progress made by EBBA1 for the quantitative assessment of species ranges, this coarse resolution limited its direct value for most practical conservation planning exercises. The higher the resolution the better the inferences that can be made concerning issues like distribution (e.g. presence in protected areas) and ecology (e.g. habitat requirements, climatic niches). Thus, in addition to the compilation of data at the 50x50 km scale, a protocol to collect standardized data across the whole of Europe was implemented in EBBA2 which allowed modelling of species occurrence at a resolution of 10x10 km. This information was mostly derived from timed visits (complete lists of species obtained at a given site and in given time) specifically designed for atlas work and from standardized common bird monitoring schemes. Given the complexity of the data, with many different sources, field methods and intensities of coverage across Europe, the EBCC convened a group of experts in a “Spatial Modelling Group” (SMOG) to approach this pan-European spatial distribution modelling in the best possible way. A final selection of more than 35 000 EBBA2 standard surveys obtained in over 17 000 different 10x10 km squares was available for modelling (Figure 3). This dataset incorporated several elements to allow the robust assessment of the detection probability for many species, i.e. repeated visits at a sample of available squares, date, field method and duration of the survey. A total of 40 environmental predictors, plus detection probability, were used to model species distribution averaging 10 different models (see Figure 4 for an example for Turtle Dove *Streptopelia turtur*). Finally, a cross-validation procedure was conducted in order to quantify the predictive accuracy of the model (Herrando *et al.* 2017).

Outlook for research

The present study documents the applications of EBBA1 in scientific publications and provides examples of why and how researchers may approach the coming EBBA2 data. This provides the possibility to forge an early connection with future research and encourage new ideas on how the new data can be used and how to increase the use of papers using this data. At the time of writing this article EBBA2 is still an ongoing project. The plan is to publish a book by late 2020 and produce a web-based product some time after publication.

We are convinced that EBBA2 will deliver data and information that will offer excellent and unique opportunities for further scientific analyses and publications. Spatio-temporal comparisons based on the atlas data are a powerful asset that could be applied to understand population dynamics of breeding birds across Europe. We see ample opportunities for use concerning the major topics that already have been addressed using the EBBA1 data, including environmental pressures according to Aichi Target B (see *Pressures on biodiversity*), such as impacts of climate and land-use changes or the spread of invasive (bird) species. Other subjects of interest could be the effects of the Natura2000 network and other conservation policies, identification of new priority areas for conservation, determination of species hotspots (also concerning other taxonomic groups), risks associated with propagation of zoonoses, and flyway research.

Within the context of biodiversity declines across Europe, the new European Breeding Bird Atlas represents a unique opportunity for researchers and policy makers to deliver valuable research to underpin conservation action across the continent. We are confident that, as it happened with its predecessor, EBBA2 will become a landmark for bird conservation in this region.

ACKNOWLEDGEMENTS

We would like to thank a lot here all national coordinators and data providers as well as ten-thousands of ornithologists and birdwatchers across Europe that built EBBA1 a few decades ago and those that are now making EBBA2 possible! We also thank the many organisations and people that provided financial support in both projects.

REFERENCES

1. **Balmer, D.E., Gillings, S., Caffrey, B.J., Swann, R.L., Downie, I.S. & Fuller, R.L.** 2013. *Bird Atlas 2007–11: the breeding and wintering birds of Britain and Ireland*. BTO Books, Thetford.
2. **Bässler, C., Müller, J., Hothorn, T., Kneib, T., Badeck, F. & Dziock, F.** 2010. Estimation of the extinction risk for high-montane species as a consequence of global warming and assessment of their suitability as cross-taxon indicators. *Ecol. Indic.* **10**: 341–352.
3. **Beck, J., Böller, M., Erhardt, A. & Schwanghart, W.** 2014. Spatial bias in the GBIF database and its effect on modelling species' geographic distributions. *Ecol. Inform.* **19**: 10–15.
4. **Bell, S., Marzano, M., Cent, J., Kobierska, H., Podjed, D., Vandzinskaite, D., Reinert, H., Armaitiene, A., Grodzińska-Jurczak, M. & Muršič, R.** 2008. What counts? Volunteers and their organisations in the recording and monitoring of biodiversity. *Biodivers. Conserv.* **17**: 3443–3454.
5. **BirdLife International.** 2004. *Birds in Europe: Population Estimates, Trends and Conservation Status*. BirdLife Conservation Series No 12.
6. **Brichetti, P. & Grattini, N.** 2008. Distribuzione, consistenza ed evoluzione delle popolazioni di basettino *Panurus biarmicus* nidificanti in Italia nel periodo 1980-2006. *Avocetta* **32**: 47–53 (in Italian).

7. **Budinski, I., Čulina, A., Mikulić, K. & Jurinović, L.** 2010. Bird species that have significantly changed breeding range on Croatian coastal area: comparison of 30 years old data and recent knowledge. *Bird Census News* **23**: 49–58.
8. **Carrascal, L.M. & Seoane, J.** 2009. Factors affecting large-scale distribution of the Bonelli's eagle *Aquila fasciata* in Spain. *Ecol. Res.* **24**: 565–573.
9. **Chiarucci, A., Bacaro, G. & Scheiner, S.M.** 2011. Old and new challenges in using species diversity for assessing biodiversity. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **366**: 2426–2437.
10. **Chiron, F., Shirley, S. & Kark, S.** 2009. Human-related processes drive the richness of exotic birds in Europe. *Proc. Royal Soc. B* **276**: 47–53.
11. **de Winter, J.C.F., Zadpoor, A.A. & Dodou, D.** 2014. The expansion of Google Scholar versus Web of Science: a longitudinal study. *Scientometrics* **98**: 1547–1565.
12. **Dickinson, J.L., Zuckerberg, B. & Bonter, D.N.** 2010. Citizen science as an ecological research tool: challenges and benefits. *Annu. Rev. Ecol. Evol. Syst.* **41**: 149–172.
13. **Dombrovski, V. C. & Ivanovski, V.V.** 2005. New data on numbers and distribution of birds of prey breeding in Belarus. *Acta Zool. Litu.* **15**: 218–227.
14. **EBBA2.** 2019. <http://ebba2.info>
15. **Eggers, J., Tröltzsch, T., Falcucci, A., Maioranow, L., Verburgz, P.H., Framstad, E., Louette, G., Maes, D., Szabolcs, N., Ozinga, W. & Delbaere, B.** 2009. Is biofuel policy harming biodiversity in Europe? *Glob. Change Biol. Bioenergy* **1**: 18–34.
16. **EuMon.** 2019. *EU-wide monitoring methods and systems of surveillance for species and habitats of community interest.* <http://eumon.ckff.si/>

- 445 17. **Flousek, J., Telenský, T., Hanzelka, J. & Reif, J.** 2015. Population trends of
446 Central European montane birds provide evidence for adverse impacts of
447 climate change on high-altitude species. *PLoS ONE* **10**: e0139465.
- 448 18. **Gaston, K.J., Jackson, S.F., Nagy, A., Cantú-Salazar, L. & Johnson, M.**
449 2008. Protected areas in Europe. *Ann. New York Acad. Sci.* **1134**: 97–119.
- 450 19. **Gaüzère, P., Jiguet, F. & Devictor, V.** 2015. Rapid adjustment of bird
451 community compositions to local climatic variations and its functional
452 consequences. *Glob. Change Biol.* **21**: 3367–3378.
- 453 20. **GBIF.** 2019. Global Biodiversity Information Facility. <http://www.gbif.org/>
- 454 21. **Greenwood, J.J.D.** 2007. Citizens, science and bird conservation. *J. Ornithol.*
455 **148**: 77–124.
- 456 22. **Greenwood, J.J.D.** 2017. The History of the EBCC Atlas of European Breeding
457 Birds. *Vogelwelt* **137**: 3–18.
- 458 23. **Hagemeijer, W. & Blair, M.** 1997. *The EBCC Atlas of European Breeding Birds.*
459 T. & A.D. Poyser, London.
- 460 24. **Harrison, J.A., Allan, D.G., Underhill, L.G., Herremans, M., Tree, A.J.,**
461 **Parker, V. & Brown, C.** 1997 *The Atlas of Southern African Birds.* BirdLife
462 South Africa, Johannesburg.
- 463 25. **Herrando, S., Keller, V., Voříšek, P., Kipson, M., Franch, M., Anton, M., Pla,**
464 **M., Villero, D., Sierdsema, H., Kampichler, C., Telenský, T., Gillings, S.,**
465 **Johnston, A., Gottschalk, T., Guélat, J., Sattler, T., Brotons, L., Titeux, N.,**
466 **Jiguet, F., Kéry, M. & Milanese, P.** 2017. High resolution maps for the new
467 European Breeding Bird Atlas: A first provision of standardised data and pilot
468 modelled maps. *Vogelwelt* **137**: 33–41.
- 469 26. **Herrando, S., Voříšek, P. & Keller, V.** 2014. The methodology of the new
470 European breeding bird atlas: finding standards across diverse situations. *Bird*
471 *Census News* **26**: 3–11.

27. **Huntley, B., Allen, J.R., Barnard, P., Collingham, Y.C. & Holliday, P.R.** 2013. Species distribution models indicate contrasting late-Quaternary histories for Southern and Northern Hemisphere bird species. *Glob. Ecol. Biogeogr.* **22**: 277–288.
28. **Huntley, B., Collingham, Y.C., Willis, S.G., & Green, R.E.** 2008. Potential impacts of climatic change on European breeding birds. *PLoS ONE*: **3**: e1439.
29. **Huntley, B., Green, R.E., Collingham, Y.C., Hill, J.K., Willis, S.G., Bartlein, P.J., Cramer, W., Hagemeijer, W.J.M. & Thomas, C.J.** 2004. The performance of models relating species geographical distributions to climate is independent of trophic level. *Ecol. Lett.* **7**: 417–426.
30. **Keil, P., Schweiger, O., Kühn, I., Kunin, W.E., Kuussaari, M., Settele, J., Henle, K., Brotons, L., Pe'er, G., Lengyel, S., Moustakas, A., Steinicke, H. & Storch, D.** 2012. Patterns of beta diversity in Europe: the role of climate, land cover and distance across scales. *J. Biogeogr.* **39**: 1473–1486.
31. **Keller, V.** 2014. EBBA2 - A New European Atlas of Breeding Birds. *Bird Census News* **26**: 3–5.
32. **Keller, V., Ayé, R., Müller, W., Spaar, R. & Zbinden, N.** 2010. Die prioritären Vogelarten der Schweiz: Revision 2010. *Ornithol. Beob.* **107**: 265–285 (in German).
33. **Koleček, J., Albrecht, T. & Reif, J.** 2014. Predictors of extinction risk of passerine birds in a Central European country. *Anim. Conserv.* **17**: 498–506.
34. **Meller, L., Thuiller, W., Pironon, S., Barbet-Massin, M., Hof, A. & Cabeza, M.** 2015. Balance between climate change mitigation benefits and land use impacts of bioenergy: conservation implications for European birds. *Glob. Change Biol. Bioenergy* **7**: 741–751.
35. **MOL.** 2019. Map of Life. <https://mol.org/>

- 498 36. **Møller, A.P., Erritzøe, J., Karadas, F. & Mousseau, T.A.** 2010. Historical
499 mutation rates predict susceptibility to radiation in Chernobyl birds. *J. Evol. Biol.*
500 **23**: 2132–2142.
- 501 37. **Møller, A.P., Garamszegi, L.Z., Peralta-Sánchez, J.M. & Soler, J.J.** 2011.
502 Migratory divides and their consequences for dispersal, population size and
503 parasite–host interactions. *J. Evol. Biol.* **24**: 1744–1755.
- 504 38. **Pauly, D. & Stergiou, K.I.** 2005. Equivalence of results from two citation
505 analyses: Thomson ISI's Citation Index and Google's Scholar service. *Ethics in*
506 *Science and Environmental Politics* **2005**: 33–35.
- 507 39. **Pedersen, P.E., Petersen, T.L., Jørgensen, P.S. & Tøttrup, A.P.** 2011. The
508 breeding population of red-backed shrike *Lanius collurio* in Gribskov,
509 Denmark. *Dansk Ornitologisk Forenings Tidsskrift* **105**: 179–182.
- 510 40. **Reif, J., Prylová, K., Šizling, A.L., Vermouzek, Z., Šťastný, K. & Bejček, V.**
511 2013. Changes in bird community composition in the Czech Republic from 1982
512 to 2004: increasing biotic homogenization, impacts of warming climate, but no
513 trend in species richness. *J. Ornithol.* **154**: 359–370.
- 514 41. **Sándor, A.D. & Domşa, C.** 2012. Special protected areas for conservation of
515 Romanian forest birds: Status assessment and possible expansion using
516 predictive tools. *Acta Zool. Bulgar.* **64**: 367–374.
- 517 42. **Secretariat of the Convention on Biological Diversity.** 2014. Global
518 Biodiversity Outlook 4. Montréal.
- 519 43. **Schmeller, D.S., Julliard, R., Bellingham, P.J., Böhm, M., Brummitt, N.,**
520 **Chiarucci, A., Couvet, D., Elmendorf, S., Forsyth, D.M., García-Moreno, J.,**
521 **Gregory, R.D., Magnusson, W.E., Martin, L.J., McGeoch, M.A., Mihoub, J.-**
522 **B., Pereira, H.M., Proença, V., van Swaay, C.A.M, Yahara, T. & Belnap, J.**
523 2015. Towards a global terrestrial species monitoring program. *J. Nat. Conserv.*
524 **25**: 51–57.

44. Sillero, N., Campos, J., Bonardi, A., Corti, C., Creemers, R., Crochet, P.-A., Isailović, J.C., Denoël, M., Ficetola, G.F., Gonçalves, J., Kuzmin, S., Lymberakis, P., de Pous, P., Rodríguez, A., Sindaco, R., Speybroeck, J., Toxopeus, B., Vieites, D.R. & Vences, M. 2014. Updated distribution and biogeography of amphibians and reptiles of Europe. *Amphibia-Reptilia* **35**: 1–31.
45. Tomiałojć, L. 2012. Changes in tree sparrow *Passer montanus* populations from urban parks. *International Studies on Sparrows* **36**: 4–17.
46. Veleviski, M., Hallmann, B., Grubač, B., Lisičanec, T., Stoyanov, E., Lisičanec, E., Avukatov, V., Božič, L. & Stumberger, B. 2010. Important bird areas in Macedonia: sites of Global and European importance. *Acrocephalus* **31**: 181–282.
47. Virkkala, R. & Rajasärkkä, A. 2007. Uneven regional distribution of protected areas in Finland: consequences for boreal forest bird populations. *Biol. Conserv.* **134**: 361–371.
48. Voříšek P. & Marchant J.H. 2003. Review of large/scale generic population monitoring schemes in Europe. *Bird Census News* **16**: 16–40.
49. Wichmann, B.A. & Hill, I.D. 1982. Algorithm AS 183: An Efficient and Portable Pseudo-Random Number Generator. *J. Royal Stat. Soc. Series C (Applied Statistics)* **31**: 188–190.
50. Wotton, S., Conway, G., Eaton, M., Henderson, I. & Grice, P. 2009. The status of the Dartford Warbler in the UK and the Channel Islands in 2006. *British Birds* **102**: 230–246.

548

549 **APPENDIX**

550 **Random selection of 100 papers that cited the *EBCC Atlas of European Breeding***
551 ***Birds* (EBBA1) that were used in this study to analyze the uses of this atlas in**
552 **research**

- 553 1. **Acquarone, C., Cucco, M. & Malacarne, G.** 2001. Short-term effects on body
554 condition and size of immunocompetent organs in the Hooded Crow. *Ital. J.*
555 *Zool.* **68**: 195–199.
- 556 2. **Ambrosini, R., Bolzern, A.M., Canova, L., Arieni, S., Møller, A.P. & Saino, N.**
557 2002. The distribution and colony size of barn swallows in relation to agricultural
558 land use. *J. Appl. Ecol.* **39**: 524–534.
- 559 3. **Arzel, C., Dessborn, L., Pöysä, H., Elmberg, J., Nummi, P. & Sjöberg, K.**
560 2014. Early springs and breeding performance in two sympatric duck species
561 with different migration strategies. *Ibis* **156**: 288–298.
- 562 4. **Backström, N. & Väli, Ü.** 2011. Sex- and species- biased gene flow in a
563 spotted eagle hybrid zone. *BMC Evol. Biol.* **11**: 100.
- 564 5. **Balčiauskienė, L., Jovaišas, A., Naruševičius, V., Petraška, A. & Skuja, S.**
565 2006. Diet of Tawny Owl (*Strix aluco*) and Long-eared Owl (*Asio otus*) in
566 Lithuania as found from pellets. *Acta Zool. Litu.* **16**: 37–45.
- 567 6. **Balmori, A., Cuesta, M.A. & Caballero, J.M.** 2002. Distribución de los
568 mosquiteros Ibérico (*Phylloscopus brehmii*) y Europeo (*Phylloscopus collybita*)
569 en los bosques de ribera de Castilla y León (España). *Ardeola* **49**: 19–27.
- 570 7. **Bässler, C., Müller, J., Hothorn, T., Kneib, T., Badeck, F. & Dziöck, F.** 2010.
571 Estimation of the extinction risk for high-montane species as a consequence of
572 global warming and assessment of their suitability as cross-taxon
573 indicators. *Ecol. Indic.* **10**: 341–352.

- 574 8. **Battisti, C., Aglitti, C., Sorace, A. & Trotta, M.** 2006. Water level decrease
575 and its effects on the breeding bird community in a remnant wetland in Central
576 Italy. *Ekologia* **25**: 252–263.
- 577 9. **Birtsas, P., Sokos, C., Papaspyropoulos, K.G., Batselas, T., Valiakos, G. &**
578 **Billinis C.** 2013. Abiotic factors and autumn migration phenology of Woodcock
579 (*Scolopax rusticola* Linnaeus, 1758, Charadriiformes: Scolopacidae) in a
580 Mediterranean area. *Ital. J. Zool.* **80**: 392–401.
- 581 10. **Bordjan, D. & Tome, D.** 2014. Rain may have more influence than temperature
582 on nest abandonment in the Great Tit *Parus major*. *Ardea* **102**: 79–86.
- 583 11. **Brichetti, P. & Grattini, N.** 2008. Distribuzione, consistenza ed evoluzione
584 delle popolazioni di basettino *Panurus biarmicus* nidificanti in Italia nel periodo
585 1980-2006. *Avocetta* **32**: 47–53 (in Italian).
- 586 12. **Burton, N.H.K., Dodd, S.G. Clark, N.A. & Ferns, P.N.** 2002. Breeding origins
587 of Redshank *Tringa totanus* wintering at two neighbouring sites on the Severn
588 Estuary: evidence for partial racial segregation. *Ringin & Migration* **21**: 19–24.
- 589 13. **Carrascal, L.M. & Seoane, J.** 2009. Factors affecting large-scale distribution of
590 the Bonelli's eagle *Aquila fasciata* in Spain. *Ecol. Res.* **24**: 565–573.
- 591 14. **Chiron, F., Shirley, S. & Kark, S.** 2009. Human-related processes drive the
592 richness of exotic birds in Europe. *Proc. R. Soc. Lond., B, Biol. Sci.* **276**: 47–53.
- 593 15. **Chmielewski, S. & Iwanczuk, C.** 2008. Liczebność ptaków szponiastych
594 Falconiformes oraz kruk *Corvus corax* w krajobrazie rolniczym pod Mogielnica
595 w roku 1981 i 2003. *Kulon* **13**: 67–75 (in Polish).
- 596 16. **Clark, H. & Sellers, R.M.** 2001. Biometrics of the Twite *Carduelis*
597 *flavirostris*. *Ringin & Migration* **20**: 328–337.
- 598 17. **Clark, J.A., Wernham, C.V., Balmer, D.E., Adams, S.Y., Griffin, B.M.,**
599 **Blackburn, J.R., Anning, D. & Milne, L.J.** 2001. Bird ringing in Britain and
600 Ireland in 1999. *Ringin & Migration* **20**: 239–288.

18. **Crochet, P.A., Lebreton, J.D. & Bonhomme, F.** 2002. Systematics of large white-headed gulls: patterns of mitochondrial DNA variation in western European taxa. *The Auk* **119**: 603–620.
19. **Delgado, M.P., Traba, J. & Morales, M.B.** 2011. Climate niche constraints in two coexisting steppe birds: the little and the great bustards. *Ardeola* **58**: 223–238.
20. **Dombrovski, V.C. & Ivanovski, V.V.** 2005. New data on numbers and distribution of birds of prey breeding in Belarus. *Acta Zool. Litu.* **15**: 218–227.
21. **Eggers, J., Tröltzsch, K., Falcucci, A., Maiorano, L., Verburg, P.H., Framstad, E., Louette, G., Maes, D., Nagy, S., Ozinga, W. & Delbaere, B.** 2009. Is biofuel policy harming biodiversity in Europe? *Glob. Change Biol. Bioenergy* **1**: 18–34.
22. **Elmberg, J., Nummi, P., Pöysä, H. & Sjöberg, K.** 2003. Breeding success of sympatric dabbling ducks in relation to population density and food resources. *Oikos* **100**: 333–341.
23. **Exo, K.M., Becker, P.H. & Sommer, U.** 1998. Umweltchemikalien in Eiern von Binnenland-und Wattenmeerbrütern des Austernfischers (*Haematopus ostralegus*). *J. Ornithol.* **139**: 401–405 (in German).
24. **Felton, A., Lindbladh, M., Elmberg, J., Felton, A.M., Andersson, E., Sekercioglu, C.H., Collingham, Y. & Huntley, B.** 2014. Projecting impacts of anthropogenic climatic change on the bird communities of southern Swedish spruce monocultures: will the species poor get poorer? *Ornis Fennica* **91**: 1–13.
25. **Finney, S.K., Wanless, S., Harris, M.P. & Monaghan, P.** 2001. The impact of gulls on puffin reproductive performance: an experimental test of two management strategies. *Biol. Conserv.* **98**: 159–165.
26. **Flousek, J., Telenský, T., Hanzelka, J. & Reif, J.** 2015. Population trends of Central European montane birds provide evidence for adverse impacts of climate change on high-altitude species. *PLoS ONE* **10**: e0139465.

27. **Fox, A.D., Ramo, C., Varo, N., Sánchez, M.I., Amat, J.A. & Green, A.J.** 2013. Late-moulting Black-necked Grebes *Podiceps nigricollis* show greater body mass in the face of failing food supply. *Ibis* **155**: 814–822.
28. **Gainzarain, J.A., Arambarri, R. & Rodríguez, A.F.** 2002. Population size and factors affecting the density of the Peregrine Falcon *Falco peregrinus* in Spain. *Ardeola* **49**: 67–74.
29. **Gaston, K. J., Jackson, S. F., Nagy, A., Cantú-Salazar, L. & Johnson, M.** 2008. Protected areas in Europe: principle and practice. *Ann. N. Y. Acad. Sci.* **1134**: 97–119.
30. **Gaüzère, P., Jiguet, F. & Devictor, V.** 2015. Rapid adjustment of bird community compositions to local climatic variations and its functional consequences. *Glob. Change Biol.* **21**: 3367–3378.
31. **Gibbons, D.W.** 2000. Development of Pan-European breeding bird monitoring. *The Ring* **22**: 25–33.
32. **Gradev, G., García, V., Ivanov, I., Zhelev, P. & Kmetova, E.** 2012. Data from Egyptian vultures *Neophron percnopterus* tagged with GPS/GSM transmitters in Bulgaria. *Acta Zool. Bulgar.* **4**: 141–146.
33. **Grussu, M.** 1999. Swamp-hen in Italy. *British Birds* **92**: 183–192.
34. **Guzmán, J.L., Ferrand, Y. & Arroyo, B.** 2011. Origin and migration of woodcock *Scolopax rusticola* wintering in Spain. *Eur. J. Wildlife Res.* **57**: 647–655.
35. **Heylen, D., Sprong, H., Oers, K., Fonville, M., Leirs, H. & Matthysen, E.** 2014. Are the specialized bird ticks, *Ixodes arboricola* and *I. frontalis*, competent vectors for *Borrelia burgdorferi* sensu lato? *Environ. Microbiol.* **16**: 1081–1089.
36. **Huntley, B., Allen, J.R., Barnard, P., Collingham, Y.C. & Holliday, P.R.** 2013. Species distribution models indicate contrasting late-Quaternary histories for

- Southern and Northern Hemisphere bird species. *Glob. Ecol. Biogeogr.* **22**: 277–288.
37. **Huntley, B., Green, R.E., Collingham, Y.C., Hill, J.K., Willis, S.G., Bartlein, P.J., Cramer, W., Hagemeljer, W.J.M. & Thomas, C.J.** 2004. The performance of models relating species geographical distributions to climate is independent of trophic level. *Ecol. Lett.* **7**: 417–426.
38. **Jenkins, R.K. & Ormerod, S.J.** 2002. Habitat preferences of breeding Water Rail *Rallus aquaticus*. *Bird Study* **49**: 2–10.
39. **Keil, P., Schweiger, O., Kühn, I., Kunin, W.E., Kuussaari, M., Settele, J., Henle, K., Brotons, L., Pe'er, G., Lengyel, S., Moustakas, A., Steinicke, H. & Storch, D.** 2012. Patterns of beta diversity in Europe: the role of climate, land cover and distance across scales. *J. Biogeogr.* **39**: 1473–1486.
40. **Keil, P., Wilson, A.M., & Jetz, W.** 2014. Uncertainty, priors, autocorrelation and disparate data in downscaling of species distributions. *Divers. Distrib.* **20**: 797–812.
41. **Keller, V.** 2000. Winter distribution and population change of Red-crested Pochard *Netta rufina* in southwestern and central Europe. *Bird Study* **47**: 176–185.
42. **Keller, V., Ayé, R., Müller, W., Spaar, R. & Zbinden, N.** 2010. Die prioritären Vogelarten der Schweiz: Revision 2010. *Ornithol. Beob.* **107**: 265–285 (in German).
43. **Kerbiriou C., Le Viol, I., Jiguet, F. & Devictor, V.** 2009. More species, fewer specialists: 100 years of changes in community composition in an island biogeographical study. *Divers. Distrib.* **15**: 641–648.
44. **Koleček, J., Albrecht, T. & Reif, J.** 2014. Predictors of extinction risk of passerine birds in a Central European country. *Anim. Conserv.* **17**: 498–506.
45. **Konter, A. & Konter, M.** 2006. Migration patterns and site fidelity of European Grebes Podicipedidae. *Regulus* **21**: 1–17.

- 684 46. **Kosinski, Z.** 2001. The breeding ecology of the Greenfinch *Carduelis chloris* in
685 urban conditions (study in Krotoszyn, W Poland). *Acta Ornithol.* **36**: 111–121.
- 686 47. **Llabrés, X.** 2010. Cria possible de pinsà trompeter *Bucanetes githagineus* a
687 Mallorca. *Anuari Ornitològic de les Balears* **25**: 51–55 (in Catalan).
- 688 48. **Lovászi, P., Bártol, I., & Moskát, C.** 2000. Nest-site selection and breeding
689 success of the Lesser Grey Shrike (*Lanius minor*) in Hungary. *The Ring* **22**:
690 157–164.
- 691 49. **Marti, J.** 2007. Zur Habitatwahl des Waldlaubsängers *Phylloscopus sibilatrix* im
692 Kanton Glarus. *Ornithol. Beob.* **104**: 45–52 (in German).
- 693 50. **Mattes, H. & Gatter, W.** 2011. Beeinflusst der Star *Sturnus vulgaris* über
694 Höhlenkonkurrenz die Häufigkeit von Spechten *Dendrocopos sp.*? *Ornithol.*
695 *Beob.* **108**: 251–259 (in German).
- 696 51. **Mazgajski, T.D.** 2002. Nesting phenology and breeding success in great
697 spotted woodpecker *Picoides major* near Warsaw (Central Poland). *Acta*
698 *Ornithol.* **37**: 1–5.
- 699 52. **Meller, L., Thuiller, W., Pironon, S., Barbet-Massin, M., Hof, A. & Cabeza, M.**
700 2015. Balance between climate change mitigation benefits and land use
701 impacts of bioenergy: conservation implications for European birds. *Glob.*
702 *Change Biol. Bioenergy* **7**: 741–751.
- 703 53. **Meltofte, H., Durinck, J., Jakobsen, B., Nordstrom, C. & Riget, F.F.** 2006.
704 Trends in wader populations in the East Atlantic flyway as shown by numbers of
705 autumn migrants in W Denmark, 1964–2003. *Wader Study Group Bulletin* **109**:
706 111–119.
- 707 54. **Mestecăneanu, A. & Mestecăneanu, F.** 2011. Researches regarding the
708 influence of the weather on the flight of the white storks (*Ciconia ciconia*) in the
709 spring migration across the Doamnei River hydrographical basin (Argeş County,
710 Romania). *Fascicula Biologie* **18**: 81–87.

- 711 55. **Mjølunes, K.R., Bunes, V. & Solbakken, K.A.** 2005. Sjeldne fugler i Norge i
712 2003. *Ornis Norvegica* **28**: 4–50 (in Norwegian).
- 713 56. **Møller, A.P., Erritzøe, J., Karadas, F. & Mousseau, T.A.** 2010. Historical
714 mutation rates predict susceptibility to radiation in Chernobyl birds. *J. Evol.*
715 *Biol.* **23**: 2132–2142.
- 716 57. **Møller, A.P., Garamszegi, L.Z., Peralta-Sánchez, J.M. & Soler, J.J.** 2011.
717 Migratory divides and their consequences for dispersal, population size and
718 parasite–host interactions. *J. Evol. Biol.* **24**: 1744–1755.
- 719 58. **Moreira, F., Ferreira, P.G., Rego, F.C. & Bunting, S.** 2001. Landscape
720 changes and breeding bird assemblages in northwestern Portugal: the role of
721 fire. *Landsc. Ecol.* **16**: 175–187.
- 722 59. **Morelli, F.** 2015. Indicator species for avian biodiversity hotspots: Combination
723 of specialists and generalists is necessary in less natural environments. *J. Nat.*
724 *Conserv.* **27**: 54–62.
- 725 60. **Müller-Buser, M.** 2000. Auftreten und Verschwinden der Sperbergrasmücke
726 *Sylvia nisoria* bei Ramosch im Unterengadin zwischen 1987 und 1999. *Ornithol.*
727 *Beob.* **97**: 233–242 (in German).
- 728 61. **Murgui, E.** 2001. Factors influencing the distribution of exotic bird species in
729 Comunidad Valenciana (Spain). *Ardeola* **48**: 149–160.
- 730 62. **Nankinov, D.N.** 1998. Wood Sandpiper *Tringa glareola* and Green Sandpiper
731 *Tringa ochropus* in Bulgaria. *International Wader Studies* **10**: 370–374.
- 732 63. **Neto, J.M. & Correia, E.** 2013. Biometrics and Sub-Specific Identification of
733 White-Spotted Bluethroats *Luscinia svecica cyanecula* and *L. s. namnetum*
734 During Autumn and Winter. *Ardeola* **59**: 309–315.
- 735 64. **Osiejuk, T.S., Olech, B., Ratyńska, K., Owsinski, A. & Gromadzka-**
736 **Ostrowska, J.** 2004. Effects of season, plasma testosterone and body size on
737 corncrake (*Crex crex*) call rhythm. *Annales Zoologici Fennici* **41**: 647–659.

- 738 65. **Palomino, D., Carrascal, L.M. & Potti, J.** 2011. Distribution of Azure-winged
739 Magpies *Cyanopica cooki* in Spain: both local and large-scale factors
740 considered. *Acta Ornithol.* **46**: 68–79.
- 741 66. **Papeş, M.** 2007. Ecological niche modeling approaches to conservation of
742 endangered and threatened birds in Central and Eastern Europe. *Biodiversity*
743 *Informatics* **4**: 14–26.
- 744 67. **Pedersen, P.E., Petersen, T.L., Jørgensen, P.S. & Tøttrup, A.P.** 2011. The
745 breeding population of red-backed shrike *Lanius collurio* in Gribskov,
746 Denmark. *Dansk Ornitologisk Forenings Tidsskrift* **105**: 179–182.
- 747 68. **Perkins, A.J., Hancock, M.H., Butcher, N. & Summers R.W.** 2005. Use of
748 time-lapse video cameras to determine causes of nest failure of Slavonian
749 Grebes *Podiceps auritus*. *Bird Study* **52**: 159–165.
- 750 69. **Piersma, T. & van Der Velde, M.** 2009. Breeding season-specific sex
751 diagnostics in the monomorphic House Martin *Delichon urbicum*. *Bird Study* **56**:
752 127–131.
- 753 70. **Polo, V., Carrascal, L.M., & Metcalfe, N.B.** 2007. The effects of latitude and
754 day length on fattening strategies of wintering coal tits *Periparus ater* (L.): a field
755 study and aviary experiment. *J. Anim. Ecol.* **76**: 866–872.
- 756 71. **Preston, C.D.** 2013. Following the BSBI's lead: the influence of the Atlas of the
757 British flora, 1962–2012. *New J. Bot.* **3**: 2–14.
- 758 72. **Procházka, P. & Reif, J.** 2002. Movements and settling patterns of Sedge
759 Warblers (*Acrocephalus schoenobaenus*) in the Czech Republic and Slovakia-
760 an analysis of ringing recoveries. *The Ring* **24**: 3–13.
- 761 73. **Qninba, A., Benhoussa, A., El Agbani, M. A., Dakki, M. & Thevenot, M.**
762 2006. Etude phénologique et variabilité interannuelle d'abondance des
763 Charadriidés (Aves, Charadrii) dans un site Ramsar du Maroc: la Merja
764 Zerga. *Bulletin de l'Institut Scientifique* **28**: 35–47 (in French).

74. **Randler, C.** 2003. Reactions to human disturbances in an urban population of the Swan Goose *Anser cygnoides* in Heidelberg (SW Germany). *Acta Ornithol.* **38**: 47–52.
75. **Reif, J. & Flousek, J.** 2012. The role of species' ecological traits in climatically driven altitudinal range shifts of central European birds. *Oikos* **121**: 1053–1060.
76. **Reif, J., Prylová, K., Šizling, A.L., Vermouzek, Z., Šťastný, K. & Bejček, V.** 2013. Changes in bird community composition in the Czech Republic from 1982 to 2004: increasing biotic homogenization, impacts of warming climate, but no trend in species richness. *J. Ornithol.* **154**: 359–370.
77. **Rönkä, N., Kvist, L., Pakanen, V.M., Rönkä, A., Degtyarev, V., Tomkovich, P., Tracy, D. & Koivula, K.** 2012. Phylogeography of the Temminck's Stint (*Calidris temminckii*): historical vicariance but little present genetic structure in a regionally endangered Palearctic wader. *Divers. Distrib.* **18**: 704–716.
78. **Salit, K. & Sol, D.** 2005. Establishment success across convergent Mediterranean ecosystems: an analysis of bird introductions. *Conserv. Biol.* **19**: 1519–1527.
79. **Sándor, A.D. & Domşa, C.** 2012. Special protected areas for conservation of Romanian forest birds: Status assessment and possible expansion using predictive tools. *Acta Zool. Bulgar.* **64**: 367–374.
80. **Schaub, M., Pradel, R., Jenni, L. & Lebreton, J.D.** 2001. Migrating birds stop over longer than usually thought: an improved capture–recapture analysis. *Ecology* **82**: 852–859.
81. **Schekkerman, H. & Visser, G.H.** 2001. Prefledging energy requirements in shorebirds: energetic implications of self-feeding precocial development. *The Auk* **118**: 944–957.
82. **Schifferli, L., Spaar, R. & Koller, A.** 2006. Fence and plough for Lapwings: Nest protection to improve nest and chick survival in Swiss farmland. *Osnabrücker Nat. Wiss. Mitt.* **32**: 123–129 (in German).

83. **Siverio, M., López-Suárez, P., Siverio, F., Rodríguez, B., Varo-Cruz, N. & López-Jurado, L.F.** 2014. Density, nest site characteristics and breeding rates of the osprey (*Pandion haliaetus*) in the southern limit of its range in the Western Palearctic (Boa Vista, Cape Verde Islands). *Afr. J. Ecol.* **52**: 50–58.
84. **Smalley, I., O'Hara-Dhand, K., McLaren, S., Svircev, Z. & Nugent, H.** 2013. Loess and bee-eaters I: ground properties affecting the nesting of European bee-eaters (*Merops apiaster* L. 1758) in loess deposits. *Quatern. Int.* **296**: 220–226.
85. **Smart, M., Azafzaf, H. & Dlensi, H.** 2007. The 'Eurasian' Spoonbill (*Platalea leucorodia*) in Africa. *Ostrich* **78**: 495–500.
86. **Sorace, A.** 2001. Value to wildlife of urban-agricultural parks: a case study from Rome urban area. *Environ. Manage.* **28**: 547–560.
87. **Staffan, B. & Hasselquist, D.** 1999. Phylogeographic population structure of great reed warblers: an analysis of mtDNA control region sequences. *Biol. J. Linnean Soc.* **66**: 171–185.
88. **Tavares, P.C., Monteiro, L.R., Lopes R.J., Correia Santos, M.M. & Furness, R.W.** 2004. Intraspecific variation of mercury contamination in chicks of black-winged stilt (*Himantopus himantopus*) in coastal wetlands from southwestern Europe. *B. Environ. Contam. Toxicol.* **72**: 437–444.
89. **Thomas, F., Deerenberg, C., Lepley, M. & Hafner, H.** 1999. Do breeding site characteristics influence breeding performance of the Purple Heron *Ardea purpurea* in the Camargue? *Rev. Ecol. (Terre Vie)* **54**: 269–281.
90. **Thorup, K.** 2004. Reverse migration as a cause of vagrancy. *Bird Study* **51**: 228–238.
91. **Tomek, T. & Guminski, W.** 2003. Bird remains from the Mesolithic and Neolithic Site Dudka Masuria NE Poland. *Acta Zool. Cracov.* **46**: 9–19.
92. **Tomiałojć, L.** 2012. Changes in tree sparrow *Passer montanus* populations from urban parks. *International Studies on Sparrows* **36**: 4–17.

93. **Velevski, M., Hallmann, B., Grubač, B., Lisičanec, T., Stoynov, E., Lisičanec, E., Avukatov, V. Božič, L. & Stumberger, B.** 2010. Important bird areas in Macedonia: sites of Global and European importance. *Acrocephalus* **31**: 181–282.
94. **Venema, P.** 1997. Pleisterende Regenwulpen *Numenius phaeopus* in Drenthe in 1980-95. *Drentse vogels* **10**: 27–36 (in German).
95. **Virkkala, R. & Rajasärkkä, A.** 2007. Uneven regional distribution of protected areas in Finland: consequences for boreal forest bird populations. *Biol. Conserv.* **134**: 361–371.
96. **Virkkala, R., Luoto, M., Heikkinen, R.K. & Leikola, N.** 2005. Distribution patterns of boreal marshland birds: modelling the relationships to land cover and climate. *J. Biogeog.* **32**: 1957–1970.
97. **Walker, D.** 2004. Audouin's Gull: new to Britain. *British Birds* **97**: 537–541.
98. **Wotton, S., Conway, G., Eaton, M., Henderson, I. & Grice, P.** 2009. The status of the Dartford Warbler in the UK and the Channel Islands in 2006. *British Birds* **102**: 230–246.
99. **Wotton, S.R., Langston, R.H. & Gregory, R.D.** 2002. The breeding status of the Ring Ouzel *Turdus torquatus* in the UK in 1999. *Bird Study* **49**: 26-34.
100. **Завьялов, Е.В., Табачишин, В.Г., Хрустов, А.В. & Якушев, Н.Н.** 2003. Современное распространение и особенности экологии журавля-красавки в саратовском Заволжье. *Беркут* **12**: 76–82 (in Russian).

842

843 LEGENDS TO FIGURES

844 **Figure 1.** 50x50 km squares for which data were reported for EBBA1 (light grey
845 squares) and for EBBA2 (dark grey dots). EBBA1 data came from 3 949 squares and
846 coverage for EBBA2 was 5 095 squares (provisional values yet). These values
847 correspond, respectively, to 74% and 96% of the total number of squares in the area
848 shown in the map. It is very important to highlight that the completeness of the
849 coverage was very low in EBBA1 in the northern half of Russia and in some parts of
850 Caucasus and the Western Balkans. Data collection for EBBA2 was mainly carried out
851 from 2013 to 2017 in the more than 50 participating countries, including much better
852 coverage in the European parts of Russia and Kazakhstan, the Caucasus countries,
853 the Canary Islands, Cyprus and the whole of Turkey.

854 **Figure 2.** Number of papers citing EBBA1 that were attributed to each of the analysed
855 main topics.


856 **Figure 3.** Location of the 17 000 10x10 km squares with EBBA2 standardized surveys.
857 In the context of this project, all these squares were surveyed in a standardized
858 manner between one and ten times. Each survey have a complete list of all species
859 recorded in a given time frame at a particular site and on a single day.

860 **Figure 4.** Pilot map showing the probability of occurrence for the Turtle Dove
861 *Streptopelia turtur* in all 10x10 km squares in Europe based on EBBA2 spatial
862 distribution modelling (light-dark grey indicates low-high probability of occurrence,
863 respectively).

864

865 FIGURES

866 Figure 1.



867

868 Figure 2.




Figure 3.




Figure 4.



873